



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Additional Notes to the 'Birds of Gallatin County, Montana'.—

These notes form an appendix to my paper published in 'The Auk,' January, 1911, pp. 26-49. The numbers are those of my list.

37. **Gallinago delicata**. WILSON'S SNIFE.— This species appears to be fairly common and regular as a winter resident in warm swamps near Bozeman. I observed seven birds on December 26, 1911.

43. **Catoptrophorus semipalmatus inornatus**. WESTERN WILLET.— Mr. W. L. Thomas of Belgrade found two nests of this species during the summer of 1911.

120. **Hesperiphona vespertina montana**. WESTERN EVENING GROSBILL.— I observed a small flock in Bozeman on January 2, 1911. The species is evidently occasionally a winter resident, though much commoner as a migrant.

126. **Leucosticte tephrocotis littoralis**. HEPBURN'S ROSY FINCH.— I observed this subspecies in company with *L. t. tephrocotis* in Bridger Cañon, March 17, 1911. Large flocks were seen and about 10 percent. of the birds were of this form.

170. **Vermivora celata lutescens**. LUTESCENT WARBLER.— I observed a male of this species in song, in an aspen grove near Bozeman, July 4, 1910. It probably breeds in the region and the breeding bird is probably *V. c. celata*, but it was neither secured nor observed closely.

199. **Cyanocephalus cyanocephalus**. PIÑON JAY.— A flock of about 40 birds was observed flying over the western part of Bozeman on Sept. 11, 1911. This adds a new species to the county list, and to my knowledge, is the most westerly record for this species in Montana.—
ARETAS A. SAUNDERS.

Two new records for Washington State.— The following birds appear to be new to the State list.

Ceryle alcyon caurina. NORTHWESTERN BELTED KINGFISHER.— A specimen taken September 17, 1896, was recently identified for me by Mr. Oberholser as this new form. It was collected by me on Puget Sound, in the vicinity of Tacoma, Wash., which I believe extends their range this much southward. As I have no other specimens, some further collecting must be done in order to prove this to be the resident form.

Zonotrichia albicollis. WHITE-THROATED SPARROW.— At Sherlock, in Thurston County, on October 13, 1912, I was so fortunate as to collect a female of this Sparrow. My bird was with a very large company of Nuttall's Sparrows (*Zonotrichia leucophrys nuttalli*), which had begun to arrive the previous evening, but a careful search failed to reveal any more *albicollis*. An interesting feature of this migration, possibly of some significance, is that the local breeding *nuttalli* had long since left for the south, and the migration in question was a considerable surprise to me. I believe my bird forms a new record for the state, if not for the Pacific coast north of Oregon.— J. HOOPER BOWLES, Tacoma, Wash.